

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



(43) International Publication Date
9 January 2003 (09.01.2003)

PCT

(10) International Publication Number
WO 03/002530 A2

(51) International Patent Classification⁷: **C07D 207/00**

NC 27709 (US). **LENHARD, James, Martin** [US/US]; GlaxoSmithKline, Five Moore Drive, Po Box 13398, Research Triangle Park, NC 27709 (US).

(21) International Application Number: PCT/US02/20466

(22) International Filing Date: 26 June 2002 (26.06.2002)

(74) Agents: **LEVY, David, J et al.**; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US).

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:

60/301,333 27 June 2001 (27.06.2001) US
60/387,011 6 June 2002 (06.06.2002) US

(81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.

(71) Applicant (*for all designated States except US*): **SMITHKLINE BEECHAM CORPORATION** [US/US]; One Franklin Plaza, Philadelphia, PA 19101 (US).

(72) Inventors; and

(75) Inventors/Applicants (*for US only*): **HAFFNER, Curt, Dale** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **MC-DOUGALD, Darryl, Lynn** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **RANDHAWA, Amarjit, Sab** [CA/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **REISTER, Steven, Michael** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park, NC 27709 (US). **DEATON, David, N.** [US/US]; GlaxoSmithKline, Five Moore Drive, PO Box 13398, Research Triangle Park,

(84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

WO 03/002530 A2

(54) Title: PYRROLIDINES AS DIPEPTIDYL PEPTIDASE INHIBITORS

(57) Abstract: The present invention relates to novel compounds, their use for inhibiting serine proteases, such as dipeptidyl peptidases, such as dipeptidyl peptidase IV (DPP-IV) and to methods for their production and their therapeutic utility.

PYRROLIDINES AS DIPEPTIDYL PEPTIDASE INHIBITORSField of Invention

The present invention relates to compounds inhibiting dipeptidyl peptidases, such as II (DPP-II) and IV (DPP-IV), to methods for their production, and to their therapeutic utility.

Background of the Invention

5 Dipeptidyl peptidase IV (DPP-IV) is a post-proline/alanine cleaving serine protease found in various tissues of the body including kidney, liver, and intestine. DPP-IV is thought to regulate the activity of multiple physiologically important peptides, including, but not limited to, GLP1, GIP, GLP2, GRP, vasoactive intestinal peptide, peptide histidine methionine, PYY, substance P, beta-casomorphine, NPY, 10 PACAP38, prolactin, chorionic gonadotropin, aprotinin, corticotropin-like intermediate lobe peptide, pituitary adenylyl cyclase-activating peptide, (Tyr)melanostatin, LD78beta(3-70), RANTES, eotaxin, procolipase, enterostatin, vasostatin 1, endomorphin, morphiceptin, stromal cell derived factor, macrophage-derived chemokine, granulocyte chemotactic protein-2, and GHRH/GRF. As examples of the therapeutic 15 value of DPP-IV, DPP-IV is believed to be involved in a variety of metabolic, gastrointestinal, viral, and inflammatory diseases, including, but not limited to, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders such as encephalomyelitis, complement mediated disorders such as glomerulonephritis, 20 lipodystrophy, and tissue damage, psychosomatic, depressive, and neuropsychiatric disease such as anxiety, depression, insomnia, schizophrenia, epilepsy, spasm, and chronic pain, HIV infection, allergies, inflammation, arthritis, transplant rejection, high

blood pressure, congestive heart failure, tumors, and stress-induced abortions, for example cytokine-mediated murine abortions. For example, DPP-IV, also known as CD26, mediates T-cell activation and HIV infection (Ohtsuki et al., 2000). T-cells expressing DPP-IV/CD26 are preferentially infected and depleted in HIV-infected individuals (Ohtsuki et al., 2000). DPP-IV inhibitors have demonstrated anti-inflammatory effects in animal models of arthritis (Tanaka et al, 1997). Additionally, DPP-IV inhibition has been shown to prolong cardiac transplant survival (Korom et al., 1997). *In vitro* studies suggest that DPP-IV/CD26 expression correlate with tumor progression of malignant melanomas of the skin (Van den Oord, 1998). Furthermore, DPP-IV is thought to regulate metabolism by cleaving the penultimate proline/alanine at the amino-terminus of polypeptides (Mentlein, 1999), such as glucagon-like peptides (GLP) and neuropeptide Y (NPY).

More specifically, GLPs help metabolize glucose and, thus, regulation of GLPs likely should be beneficial in the treatment of metabolic disorders such as diabetes. Diabetes, for example type 2 (also called noninsulin-dependent diabetes mellitus (NIDDM) or maturity-onset) diabetes, results in elevated blood sugar levels due to absolute or relative insufficiencies of insulin. Type 2 diabetes is the more common form of diabetes, accounting for 90% of cases, or about 16 million Americans. Most type 2 diabetics produce variable, sometimes normal, amounts of insulin, but they have abnormalities in liver and muscle cells that resist its actions. Insulin attaches to the receptors of cells, but glucose does not get inside, a condition known as insulin resistance. Many type 2 diabetics seem to be incapable of secreting enough insulin to overcome insulin resistance. GLP-1 enhances insulin secretion. Thus, regulation of GLP-1 correlates to a regulation of insulin secretion. Moreover, GLP-1 decreases

hepatic glucose production, gastric emptying, and food intake (Deacon et al., 1995). Further, GLP-2 maintains the integrity of the intestinal mucosal epithelium via effects on gastric motility, nutrient absorption, crypt cell proliferation and apoptosis, and intestinal permeability (Drucker, 2001).

5 DPP-IV inhibitors preserve GLP-1 function for a longer time (Balka, 1999). Thus, DPP-IV inhibitors may promote satiety, weight loss, and the antidiabetic effects of GLP-1 (Deacon et al., 1995; Holst and Deacon, 1998). For example, inhibition of DPP-IV with the known compound NVP-DPP728 increases plasma GLP-1 (2-36 amide) concentrations and improves oral glucose tolerance in obese Zucker rats. See,
10 Diabetologia 42: 1324-1331. Both subcutaneously and intravenously administered GLP-1 is rapidly degraded from the NH₂-terminus in type II diabetic patients and in healthy subjects. See, Diabetes 44:1126, 1995.

 Moreover, DPP-IV inhibitors preserve GLP-2 for longer periods of time and, thus, may be useful for treating intestinal insufficiencies and mucous membrane
15 disorders (Hartmann B et al., 2000).

 While DPP-IV is the predominate protease regulating GLP turnover, similar substrate or inhibitor specificity may be observed for related proteases. Related serine proteases include, but are not limited to, dipeptidyl peptidase-II (DPP-II), dipeptidyl peptidase IV beta, dipeptidyl peptidase 8, dipeptidyl peptidase 9, aminopeptidase P,
20 fibroblast activating protein alpha (seprase), prolyl tripeptidyl peptidase, prolyl oligopeptidase (endoproteinase Pro-C), attractin (soluble dipeptidyl-aminopeptidase), acylaminoacyl-peptidase (N-acylpeptide hydrolase; fMet aminopeptidase) and lysosomal Pro-X carboxypeptidase (angiotensinase C, prolyl carboxypeptidase). Proline-cleaving metallopeptidases that may share similar substrate or inhibitor

specificity to DPP-IV include membrane Pro-X carboxypeptidase (carboxypeptidase P), angiotensin-converting enzyme (Peptidyl-dipeptidase A multi-peptidase), collagenase 1 (interstitial collagenase; matrix metalloproteinase 1; MMP-1; Mcol-A), ADAM 10 (alpha-secretase, myelin-associated disintegrin metalloproteinase), neprilysin (atriopeptidase; CALLA; CD10; endopeptidase 24.11; enkephalinase), Macrophage elastase (metalloelastase; matrix metalloproteinase 12; MMP-12), Matrilysin (matrix metalloproteinase 7; MMP-7), and neurolysin (endopeptidase 24.16; microsomal endopeptidase; mitochondrial oligopeptidase). See <http://merops.iapc.bbsrc.ac.uk/>.

Furthermore, beyond mammalian serine peptidases and proline-cleaving metallopeptidases, other non-mammalian proteases may share similar substrate or inhibitor specificity to DPP-IV. Non-limiting examples of such non-mammalian serine proteases include prolyl aminopeptidase (prolyl iminopeptidase), IgA1-specific serine type prolyl endopeptidase (IgA protease, *Neisseria*, *Haemophilus*), dipeptidyl aminopeptidase A (STE13) (*Saccharomyces cerevisiae*), dipeptidyl aminopeptidase B (fungus), prolyl oligopeptidase homologue (*Pyrococcus* sp.), oligopeptidase B (*Escherichia coli* alkaline proteinase II; protease II), dipeptidyl aminopeptidase B1 (*Pseudomonas* sp.), dipeptidyl-peptidase IV (bacteria), dipeptidyl aminopeptidase (*Aureobacterium*), dipeptidyl-peptidase IV (insect), dipeptidyl-peptidase V, allergen Tri t 4 (*Trichophyton tonsurans*), secreted alanyl DPP (*Aspergillus oryzae*), peptidase II-mes (*Prosopis velutina*), and bamboo serine proteinase (*Pleioblastus hindsii*). Non-limiting examples of such non-mammalian proline-cleaving metallopeptidases include penicillolysin (fungal acid metalloendopeptidase), proline-specific peptidyl-dipeptidase (*Streptomyces*), coccolysin (gelatinase, *Enterococcus faecalis*), aminopeptidase Ey, (hen egg yolk) (apdE g.p.; *Gallus gallus domesticus*), gametolysin (*Chlamydomonas*


cell wall degrading protease), and snake venom proline-cleaving metalloproteases as well. See <http://merops.iapc.bbsrc.ac.uk/> for further reference.

Dipeptidyl peptidase II (DPP II) is a serine protease localized to lysosomes in cells and believed to be involved in lysosomal degradation and protein turnover. The order of expression of DPP-II is kidney >> testis > or = heart > brain > or = lung > spleen > skeletal muscle > or = liver (Araki H et al., J Biochem (Tokyo) 2001, 129:279-88). This expression suggests possible utility in kidney or lysosomal-related disorders. Substrate specificity studies indicated that purified DPP-II hydrolyzes specifically alanine or proline residues at acidic pH (4.5-5.5). DPP-II has significant sequence
10 homology and substrate specificity to quiescent cell proline dipeptidase and prolyl carboxypeptidase, suggesting possible overlapping functions between these proteases (Araki H et al., J Biochem (Tokyo) 2001, 129:279-88).


The present invention includes novel DPP-II and/or DPP-IV inhibitors, as well as methods of their therapeutic use and methods of their production. While not being
15 limited thereby, the compounds of the present invention are believed useful for the treatment of a variety of metabolic, gastrointestinal, viral, and inflammatory diseases, including, but not limited to, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders such as encephalomyelitis, complement mediated disorders such as
20 glomerulonephritis, lipodystrophy, and tissue damage, psychosomatic, depressive, and neuropsychiatric disease such as anxiety, depression, insomnia, schizophrenia, epilepsy, spasm, and chronic pain, HIV infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions, for example cytokine-mediated murine abortions.

SUMMARY OF THE INVENTION

The present invention includes compounds of formula (I):




- including salts, solvates, and pharmaceutically functional derivatives thereof, wherein
- 5 X is $-S(O)_b-$ or $-CH_2-$; b is 0-2; R^1 and R^2 each are either H; alkyl; optionally substituted aryl or heteroaryl; or combine to form a 3 to 14 membered ring system, optionally containing one or more heteroatoms, and optionally containing one or more degrees of unsaturation; when R^1 and R^2 are optionally substituted aryl or heteroaryl, then R^3 is H or alkyl; when R^1 and R^2 are otherwise, then R^3 is



- where n is 0-5; m is 0 to 12; Y is $S(O)_p$, O, alkylene, alkenylene, alkynylene, or a bond; p is 0-2; R^4 is R^5 when Y is S, O, alkylene, alkenylene, alkynylene, or a bond, where R^5 is alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, or heterocyclyl; and R^4 is R^6 when Y is $S(O)$ or $S(O)_2$, where R^6 is alkyl, aryl, cycloalkyl, heteroaryl, amino, alkylamino,
- 15 arylamino, heteroarylamino, or cycloalkylamino.

In one embodiment, R^1 and R^2 are each alkyl and R^3 is



Preferably, each alkyl is C_1 - C_6 alkyl. More preferably, each of alkyl is methyl.

In a particular compound included within the present invention X is $-S(O)_b-$, b is 0, n is 0, m is 1, Y is $-S(O)_p-$, p is 0, R^4 is R^5 , and R^5 is optionally substituted aryl.

Preferably, R^5 is phenyl substituted with alkoxy.

In a particular compound included within the present invention X is $-CH_2-$, n is 0, m is 1, Y is $-S(O)_p-$, p is 0, R^4 is R^5 , and R^5 is optionally substituted aryl. Preferably, R^5 is phenyl substituted with alkoxy.

In a particular compound included within the present invention X is $-CH_2-$, n is 0, m is 1, Y is $S(O)_p$, p is 2, R^4 is R^6 , and R^6 is optionally substituted aryl. Preferably, R^6 is phenyl substituted with alkoxy.

In a particular compound included within the present invention X is $S(O)_b$, b is 2, n is 0, m is 1, Y is $S(O)_p$, p is 2, R^4 is R^6 , and R^6 is optionally substituted aryl. Preferably, R^6 is phenyl substituted with alkoxy.

In certain embodiments, preferably the depicted NH_2 group is *cis* to the depicted nitrile warhead. In other embodiments, preferably the depicted NH_2 group is *trans* to the depicted nitrile warhead.

Particularly preferred compounds of the present invention include:

- (4*R*)-3-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl}-1,3-thiazolidine-4-carbonitrile hydrochloride;
- (2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl}pyrrolidine-2-carbonitrile hydrochloride;
- (2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl}pyrrolidine-2-carbonitrile hydrochloride;
- (4*R*)-3-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl}-1,3-thiazolidine-4-carbonitrile 1,1-dioxide hydrochloride; and
- (2*S*)-1-[(2*S*)-2-Amino-3,3-bis(4-fluorophenyl)propanoyl]pyrrolidine-2-carbonitrile hydrochloride.

The present invention also includes pharmaceutical formulation comprising a compound of the present invention as herein described. More preferably, the pharmaceutical formulation further includes a pharmaceutically acceptable carrier.

The present invention also includes a method of inhibiting a post
5 proline/analine cleaving protease comprising administering a compound of the present invention as herein described. Preferably, the post proline/analine cleaving protease is a serine protease. Preferably, the serine protease is a dipeptidyl peptidase. In one aspect preferably the dipeptidyl peptidase is DPP-II. In another aspect preferably the dipeptidyl peptidase is DPP-IV.

10 The present invention also includes a method for the treatment or prophylaxis of metabolic disorders, gastrointestinal disorders, viral disorders, inflammatory disorders, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders, encephalomyelitis, complement mediated disorders, glomerulonephritis, lipodystrophy,
15 tissue damage, psychosomatic, depressive, and neuropsychiatric disorders, HIV infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions comprising administering a compound of the present invention as herein described. Preferably, the compound of the present invention as herein described is administered for the
20 treatment or prophylaxis of diabetes, more preferably Type II diabetes.

The present invention also includes the use of a compound of the present invention as herein described in the manufacture of a medicament for the inhibition of a post proline/analine cleaving protease. As mentioned, preferably the post proline/analine cleaving protease is a serine protease. More preferably, the serine

protease is a dipeptidyl peptidase. In one aspect preferably the dipeptidyl peptidase is DPP-II. In another aspect preferably the dipeptidyl peptidase is DPP-IV.

The present invention also includes the use of a compound of the present invention as herein described in the manufacture of a medicament for the treatment
5 or prophylaxis of metabolic disorders, gastrointestinal disorders, viral disorders, inflammatory disorders, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders, encephalomyelitis, complement mediated disorders, glomerulonephritis, lipodystrophy, tissue damage, psychosomatic, depressive, and neuropsychiatric disorders, HIV
10 infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions.

The present invention also includes a compound the present invention as herein described for use as an active therapeutic substance. Additionally, the present invention includes a compound the present invention as herein described for use in
15 the manufacture of a medicament for the inhibition of serine protease. Further, the present invention includes a compound the present invention as herein described for use in the manufacture of a medicament for the treatment or prophylaxis of metabolic disorders, gastrointestinal disorders, viral disorders, inflammatory disorders, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders,
20 psoriasis, intestinal distress, constipation, autoimmune disorders, encephalomyelitis, complement mediated disorders, glomerulonephritis, lipodystrophy, tissue damage, psychosomatic, depressive, and neuropsychiatric disorders, HIV infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions.

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENT

The term "alkyl" refers to a straight or branched chain saturated aliphatic hydrocarbon that may be optionally substituted, with multiple degrees of substitution being allowed. Examples of "alkyl" include, but are not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, t-butyl, n-pentyl, isobutyl, and the like.

As used throughout this specification, the preferred number of carbon atoms will be represented by, for example, the phrase "C_x-C_y alkyl" which refers to an alkyl group, as herein defined, containing the specified number of carbon atoms. Similar terminology will apply for other preferred ranges as well.

10 The term "alkylene" refers to a divalent straight or branched chain aliphatic hydrocarbon radical that may be optionally substituted, with multiple degrees of substitution being allowed. An example of "alkylene" includes, without limitation, methylene, namely -CH₂-.

The term "alkenyl" refers to a straight or branched chain aliphatic hydrocarbon, containing one or more carbon-to-carbon double bonds that may be optionally substituted, with multiple degrees of substitution being allowed. Examples include, but are not limited to, vinyl and the like.

As used herein the term "alkenylene" refers to a divalent straight or branched chain aliphatic hydrocarbon radical, containing one or more carbon-to-carbon double bonds, which may be optionally substituted, with multiple degrees of substitution being allowed. An example of "alkenylene" includes, without limitation, vinylene, namely, -CH=CH-.

As used herein the term "alkynyl" refers to a straight or branched aliphatic hydrocarbon containing one or more triple bond, which may optionally be substituted,

with multiple degrees of substitution being allowed. Examples of "alkynyl" as used herein include, but are not limited to, ethynyl and the like.


As used herein the term "alkynylene" refers to a divalent straight or branched chain aliphatic hydrocarbon radical, containing at least one carbon-to-carbon triple bond, that may be further substituted, with multiple degrees of substitution being allowed. An example of "alkynylene" includes, without limitation, ethynylene, namely $-C\equiv C-$.

The term "aryl" refers to an aromatic ring system, such as an optionally substituted benzene ring system, such as phenyl. The term encompasses fused systems where one or more optionally substituted benzene rings form, for example, anthracene, phenanthrene, or naphthalene ring systems. The term includes ring(s) optionally substituted, with multiple degrees of substitution being allowed, and also includes an optional alkylene linker, such as C_1-C_6 alkylene, through which the aryl group may be attached. Examples of "aryl" groups include, but are not limited to phenyl, benzyl, 2-naphthyl, 1-naphthyl, biphenyl, as well as substituted derivatives thereof.


The term "heteroaryl" refers to a monocyclic aromatic ring system, or to a fused bicyclic aromatic ring system comprising two or more aromatic rings. These heteroaryl rings contain one or more nitrogen, sulfur, and/or oxygen atoms, where N-oxides, sulfur oxides, and dioxides are permissible heteroatom substitutions and the rings may be optionally substituted, with multiple degrees of substitution being allowed. The term includes ring(s) optionally substituted, with multiple degrees of substitution being allowed, and also includes an optional alkylene linker, such as C_1-C_6 alkylene, through which the heteroaryl group may be attached. Examples of

"heteroaryl" groups used herein include furan, thiophene, pyrrole, imidazole, pyrazole, triazole, tetrazole, thiazole, oxazole, isoxazole, oxadiazole, thiadiazole, isothiazole, pyridine, pyridazine, pyrazine, pyrimidine, quinoline, isoquinoline, benzofuran, benzothiophene, indole, indazole, and substituted versions thereof.

- 5 As used herein, the term "cycloalkyl" refers to a mono- or bi-cyclic hydrocarbon ring system, which may be further substituted with multiple degrees of substitution being allowed, and which optionally includes an alkylene linker through which the cycloalkyl may be attached. Exemplary "cycloalkyl" groups include, but are not limited to, cyclopropyl, cyclobutyl, cyclopentyl, cyclohexyl, and cycloheptyl. When
- 10 substituted, one preferred substituent location for cycloalkyl groups of the present invention is at the "1-position." To illustrate, without limitation, a preferred location for a substituent is represented below with the substituent referred to as "R":



- The term "cycloalkyl" includes bridged or fused ring systems, as well, such as
- 15 hydrindane, decalin, or adamantyl. For ease of reference, also included within the term are cycloalkyl/aryl fused systems where, for example, a cycloalkyl, such as cyclohexyl, is fused with an aromatic ring, such as a benzene ring, to form groups such
- as



As used herein, the term "heterocyclic" or the term "heterocyclyl" refers to a heterocyclic ring, preferably three to twelve-membered, that is either saturated or has one or more degrees of unsaturation. These heterocyclic rings contain one or more heteroatom, such as nitrogen, sulfur, and/or oxygen atoms, where N-oxides, sulfur
5 oxides, and dioxides are permissible heteroatom substitutions. As used herein heterocyclic groups optionally may be substituted, with multiple degrees of substitution being allowed, and also includes an optional alkylene linker, such as C₁-C₆ alkylene, through which the heterocyclyl group may be attached. Such a ring may be optionally fused to one or more of another "heterocyclic" ring(s), aryl ring(s), or
10 cycloalkyl ring(s). Examples of "heterocyclic" include, but are not limited to, tetrahydrofuran, pyran, 1,4-dioxane, 1,3-dioxane, piperidine, pyrrolidine, morpholine, tetrahydrothiopyran, tetrahydrothiophene, and the like.

The term "halogen" refers to fluorine, chlorine, bromine, or iodine.

As used herein, the term "alkoxy" refers to the group -OR_a, where R_a is alkyl as
15 herein defined.

As used herein, the term "amino" refers to the group -NH₂.

As used herein, the term "alkylamino" refers to the group -N(R_a)₂, where one R_a is alkyl and the other R_a independently is H or alkyl, as herein defined.

As used herein, the term "cycloalkylamino" refers to the group -N(R_a)₂, where
20 one R_a is cycloalkyl and the other R_a independently is H or cycloalkyl, as herein defined.

As used herein, the term "arylamino" refers to the group -N(R_a)₂, where one R_a is aryl and the other R_a independently is H or aryl, as herein defined.

As used herein, the term "heteroarylamino" refers to the group $-N(R_a)_2$, where one R_a is heteroaryl and the other R_a independently is H or heteroaryl, as herein defined.

Also, as used herein throughout the present specification, the phrase

5 "optionally substituted" denotes an optional substitution, one or more times, with acyl; alkyl; alkenyl; alkynyl; alkylsulfonyl; alkoxy; cyano; halogen; haloalkyl; hydroxy; nitro; aryl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; heteroaryl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano,

10 halogen, haloalkyl, hydroxy, or nitro; arylsulfonyl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; heteroarylsulfonyl, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; aryloxy, which may be further substituted with acyl, alkoxy, alkyl, alkenyl,

15 alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; heteroaryloxy, which may be further substituted with acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro; $-R'OR^4$; or $-NR^4R^5$; where for each occurrence R' is alkylene, alkenylene, or alkynylene, and R^4 and R^5 are each independently selected from H, alkyl, alkenyl, alkynyl, cycloalkyl, heterocyclyl, aryl,

20 heteroaryl, alkylsulfonyl, arylsulfonyl, or heteroarylsulfonyl, where each occurrence of such aryl or heteroaryl may be substituted with one or more acyl, alkoxy, alkyl, alkenyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro, or R^4 and R^5 may combine to form a ring, optionally having additional heteroatoms, optionally having one or more degrees of unsaturation, and optionally being further substituted with

acyl, alkoxy, alkyl, alkenyl, alkynyl, alkylsulfonyl, cyano, halogen, haloalkyl, hydroxy, or nitro.

The compounds of the present invention may have the ability to crystallize in more than one form, a characteristic known as polymorphism. All polymorphic forms ("polymorphs") are within the scope of the present invention. Polymorphism generally can occur as a response to changes in temperature or pressure, or both, and can also result from variations in the crystallization process. Polymorphs can be distinguished by various physical characteristics that are known in the art such as x-ray diffraction patterns, solubility, and melting point.

Certain of the compounds described herein contain one or more chiral centers, or may otherwise be capable of existing as multiple stereoisomers. The scope of the present invention includes pure stereoisomers as well as mixtures of stereoisomers, such as purified enantiomers/diastereomers or enantiomerically/diastereomerically enriched mixtures. Also included within the scope of the invention are the individual isomers of the compounds *per se*, as well as any wholly or partially equilibrated mixtures thereof. The present invention covers the individual isomers of the compounds represented by the formulas above as mixtures with isomers thereof in which one or more chiral centers are inverted.

As noted above, the present invention includes salts, solvates, and pharmaceutically functional derivatives of the compounds of the present invention. Salts include addition salts, metal salts, or optionally alkylated ammonium salts. Examples of such salts include hydrochloric, hydrobromic, hydroiodic, phosphoric, sulfuric, trifluoroacetic, trichloroacetic, oxalic, maleic, pyruvic, malonic, succinic, citric, mandelic, benzoic, cinnamic, methane sulphonic, ethane sulphonic, picric, and the like.

Further salts include lithium, sodium, potassium, magnesium, and the like. Still further salts include acetate, benzenesulfonate, benzoate, bicarbonate, bisulfate, bitartrate, borate, bromide, calcium edetate, camsylate, carbonate, chloride, clavulanate, citrate, dihydrochloride, edetate, edisylate, estolate, esylate, fumarate, gluceptate, gluconate, glutamate, glycolylarsanilate, hexylresorcinate, hydrabamine, hydroxynaphthoate, isethionate, lactate, lactobionate, laurate, malate, mandelate, mesylate, methylbromide, methylnitrate, methylsulfate, monopotassium maleate, mucate, napsylate, nitrate, N-methylglucamine, pamoate (embonate), palmitate, pantothenate, phosphate/diphosphate, polygalacturonate, potassium, salicylate, sodium, stearate, subacetate, tannate, tartrate, teoclate, tosylate, triethiodide, trimethylammonium, and valerate salts. Reference is also made to *Journal of Pharmaceutical Science*, 1997, 66, 2, incorporated herein by reference, as relevant to salts.

As used herein, the term "solvate" refers to a complex of variable stoichiometry formed by a solute or a salt or pharmaceutically functional derivative thereof and a solvent. Such solvents for the purpose of the invention should not interfere with the biological activity of the solute. Examples of solvents include, but are not limited to water, methanol, ethanol, and acetic acid. Preferably the solvent used is a pharmaceutically acceptable solvent. Examples of pharmaceutically acceptable solvents include water, ethanol, and acetic acid.

The term "pharmaceutically functional derivative" refers to any pharmaceutically acceptable derivative of a compound of the present invention, for example, an ester or an amide, which upon administration to a mammal is capable of providing (directly or indirectly) a compound of the present invention or an active metabolite or residue thereof. Such derivatives are recognizable to those skilled in

the art, without undue experimentation. Nevertheless reference is made to the teaching of *Burger's Medicinal Chemistry and Drug Discovery*, 5th Edition, Vol 1: Principles and Practice, which is incorporated herein by reference to the extent of teaching pharmaceutically functional derivatives.

- 5 While compounds of the present invention may be administered as the raw chemical, preferably the compounds of the present invention are presented as an active ingredient within a pharmaceutical formulation as known in the art. Accordingly, the present invention further includes a pharmaceutical formulation comprising a compound of the present invention, or salt, solvate, or pharmaceutically
- 10 functional derivative thereof together with one or more pharmaceutically acceptable carriers. Optionally, other therapeutic and/or prophylactic ("active") ingredients may be included in the pharmaceutical formulation as well. For example, the compounds of the present invention may be combined with other anti-diabetic agents, such as one or more of the following agents: insulin, α -glucosidase inhibitors, biguanides,
- 15 insulin secretagogue, or insulin sensitizers. Non-limiting examples of α -glucosidase inhibitors include acarbose, emiglitate, miglitol, and voglibose. Non-limiting examples of biguanides include metformin, buformin, and phenformin. Non-limiting examples of insulin secretagogues include sulphonylureas. Non-limiting examples of insulin sensitizers include peroxisome proliferator activated receptor (PPAR) ligands, such as
- 20 PPAR- γ agonists, for example Actos™ and Avandia™.

Formulations of the present invention include those especially formulated for oral, buccal, parental, transdermal, inhalation, intranasal, transmucosal, implant, or rectal administration. Among the variety of administrations, oral administration typically is preferred. For oral administration tablets, capsules, and caplets may

contain conventional excipients such as binding agents, fillers, lubricants, disintegrants, and/or wetting agents. Non-limiting examples of binding agents include syrup, acacia, gelatin, sorbitol, tragacanth, mucilage of starch, or polyvinylpyrrolidone (PVP). Non-limiting examples of fillers include, for example, 5 lactose, sugar, microcrystalline cellulose, maize-starch, calcium phosphate or sorbitol. Non-limiting examples of lubricants include, for example, magnesium stearate, stearic acid, talc, polyethylene glycol or silica. Non-limiting examples of disintegrants include, for example, potato starch or sodium starch glycollate. A non-limiting example of a wetting agent includes sodium lauryl sulfate. The tablets additionally 10 may be coated according to methods known in the art.

Alternatively, the compounds of the present invention may be incorporated into oral liquid preparations such as aqueous or oily suspensions, solutions, emulsions, syrups, or elixirs. Moreover, formulations containing these compounds may be presented as a dry product for constitution with water or other suitable vehicle before 15 use. Liquid preparations may contain conventional additives. Non-limiting examples of such additives include suspending agents such as sorbitol syrup, methyl cellulose, glucose/sugar syrup, gelatin, hydroxyethylcellulose, carboxymethyl cellulose, aluminum stearate gel or hydrogenated edible fats. Additionally, emulsifying agents such as lecithin, sorbitan mono-oleate or acacia; non-aqueous vehicles (which may 20 include edible oils) such as almond oil, fractionated coconut oil, oily esters, propylene glycol or ethyl alcohol may be included. Further, preservatives such as methyl or propyl p-hydroxybenzoates or sorbic acid, may be incorporated into the preparation. Such preparations may also be formulated as suppositories, for example, containing conventional suppository bases such as cocoa butter or other glycerides.

Additionally, formulations of the present invention may be formulated for parenteral administration by injection or continuous infusion. Formulations for injection may take such forms as suspensions, solutions, or emulsions in oily or aqueous vehicles, and may contain formulatory agents such as suspending, stabilizing and/or dispersing agents. Alternatively, the active ingredient may be in powder form for constitution with a suitable vehicle, for example, sterile, pyrogen-free water, before use.

The formulations according to the invention may also be formulated as a depot preparation. Such long acting formulations may be administered by implantation, for example, subcutaneously or intramuscularly, or by intramuscular injection. Accordingly, the compounds of the invention may be formulated with suitable polymeric or hydrophobic materials, such as an emulsion in an acceptable oil, ion exchange resins, or as sparingly soluble derivatives, such as a sparingly soluble salt.

Pharmaceutical formulations may be presented in unit dose forms containing a predetermined amount of active ingredient per unit dose. Such a unit may contain certain amounts of a compound of the present invention depending on the condition being treated, the route of administration, and the age, weight and condition of the patient. Examples of such amounts include the formulation containing about 0.1 to about 99.9% active ingredient. Preferred unit dosage formulations are those containing a predetermined dose, such as a daily dose, or an appropriate fraction thereof, of an active ingredient. Such pharmaceutical formulations may be prepared by any of the methods well known in the pharmacy art.

As used herein, the term "effective amount" means that amount of a drug or pharmaceutical agent that will elicit the biological or medical response of a tissue,

system, animal, or human that is being sought, for instance, by a researcher or clinician. Furthermore, the term "therapeutically effective amount" means any amount which, as compared to a corresponding subject who has not received such amount, results in improved treatment, healing, prevention, or amelioration of a
5 disease, disorder, or side effect, or a decrease in the rate of advancement of a disease or disorder. The term also includes within its scope amounts effective to enhance normal physiological function.

The therapeutically effective amount of a compound of the present invention will depend upon a number of factors including, for example, the age and weight of
10 the animal, the precise condition requiring treatment and its severity, the nature of the formulation, and the route of administration. Therapeutic effectiveness ultimately will be at the discretion of the attendant physician or veterinarian. An effective amount of a salt or solvate, or pharmaceutically functional derivative thereof, may be determined as a proportion of the effective amount of a compound of the present
15 invention *per se*. Dosages may vary, depending upon the appropriate inhibition of DPP-IV for purposes of treatment or prophylaxis of a variety of metabolic, gastrointestinal, viral, and inflammatory diseases, including, but not limited to, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders such as
20 encephalomyelitis, complement mediated disorders such as glomerulonephritis, lipodystrophy, and tissue damage, HIV infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions, for example cytokine-mediated murine abortions.

No toxicological effects are indicated/expected when a compound of the present invention is administered in the above mentioned dosage range.

The present invention should be interpreted to cover all combinations of particular and preferred groups herein described. The application of which this
5 description and claims forms part may be used as a basis for priority in respect of any subsequent application. The claims of such subsequent application may be directed to any feature or combination of features described herein. They may take the form of product, composition, process, or use claims and may include, by way of example and without limitation, the claims hereto appended.

10 The following examples illustrate aspects of this invention, but should not be construed as limitations. As used herein the symbols and conventions used in these processes, schemes and examples are consistent with those used in the contemporary scientific literature, for example, the *Journal of the American Chemical Society* or the *Journal of Biological Chemistry*. Unless otherwise noted, all starting materials were
15 obtained from commercial suppliers and used without further purification.

¹H NMR spectra were recorded on a Varian VXR-300, a Varian Unity-300, a Varian Unity-400 instrument, or a General Electric QE-300. Chemical shifts are expressed in parts per million (ppm, δ units). Coupling constants are in units of hertz (Hz). Splitting patterns describe apparent multiplicities and are designated as s
20 (singlet), d (doublet), t (triplet), q (quartet), m (multiplet), br (broad).


Low-resolution mass spectra (MS) were recorded on a JOEL JMS-AX505HA, JOEL SX-102, or a SCIEX-APIiii spectrometer; high resolution MS were obtained using a JOEL SX-102A spectrometer. All mass spectra were taken under electrospray ionization (ESI), chemical ionization (CI), electron impact (EI) or by fast atom

bombardment (FAB) methods. Infrared (IR) spectra were obtained on a Nicolet 510 FT-IR spectrometer using a 1-mm NaCl cell. All reactions were monitored by thin-layer chromatography on 0.25 mm E. Merck silica gel plates (60F-254), visualized with UV light, 5% ethanolic phosphomolybdic acid or p-anisaldehyde solution. Flash column chromatography was performed on silica gel (230-400 mesh, Merck). Optical rotations were obtained using a Perkin Elmer Model 241 Polarimeter. Melting points were determined using a Mel-Temp II apparatus and are uncorrected.

IUPAC names are included to further identify particular compounds of the present invention. The IUPAC names stated herein should in no way limit the scope of the present invention.

EXPERIMENTALS

In accordance with the present invention and as below, one embodiment of the compounds of the present invention can be prepared by reacting a compound of formula II with an α -amino carboxylate or with an α -amino activated carboxylate, both designated herein generally as aminocarboxylates, under standard coupling conditions, for example, with HATU, DMF, Hunigs base.



More specifically, a compound of formula II may be reacted with an amino carboxylate, where the amino carboxylate is suitably protected, for example on the α -


nitrogen, with an appropriate protecting group such as, for example, a t-butyl carboxy protecting group.

In an alternate embodiment, a compound of formula II may be reacted with an amino activated carboxylate, such as, for example, N-hydroxysuccinimide ester or acid chloride, where the amino activated carboxylate is suitably protected, for example, on the α -nitrogen with an appropriate protecting group such as, for example, a t-butyl carboxy protecting group. Removal of the protecting group under suitable conditions, such as, for example, trifluoroacetic acid for the removal of the t-butyl carboxy, then generates compounds of formula (I).

For further detail regarding the preparation of amino carboxylates for use in preparing the compounds of the present invention, reference may be had to WO 95/15309 and WO 98/19998, each herein incorporated by reference as related to the preparation of such reactants.

Examples

15 Example 1



(4R)-3-[(2R)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl]-1,3-thiazolidine-4-carbonitrile hydrochloride

A. *tert*-Butyl (1*R*)-1-[[[(4*R*)-4-(aminocarbonyl)-1,3-thiazolidin-3-yl]carbonyl]-2-[(4-methoxybenzyl)thio]-2-methylpropylcarbamate.

To a stirred solution of *N*-BOC-L-Pen(MOB)-OH (300 mg, 0.812 mmol) in DMF (8 mL) was added (4*R*)-1,3-thiazolidine-4-carboxamide hydrochloride (149 mg, 0.812 mmol), HATU (309 mg, 0.812 mmol), and *N,N*-diisopropylethylamine (0.424 mL, 2.44 mmol). The reaction mixture was stirred at RT for 16 hours. Water (8 mL) was added and the reaction mixture was extracted with five portions of EtOAc. The combined extracts were washed with water, saturated CuSO₄, brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The crude material was flash chromatographed over silica gel (1:1 hexanes:EtOAc) to afford 288 mg (74% yield) of compound A as a colorless oil.

¹H NMR (CDCl₃) 400 MHz δ 7.18 (d, *J*=8.6 Hz, 2H), 6.9 (br s, 1H), 6.83 (d, *J*=8.5 Hz, 2H), 5.52 (br s, 1H), 5.41 (d, *J*=9.4 Hz, 1H), 5.10 (dd, *J*=6.9, 2.8 Hz, 1H), 4.87 (d, *J*=8.4 Hz, 1H), 4.70 (m, 2H), 3.81 (s, 2H), 3.78 (s, 3H), 3.43 (d, *J*=14.1 Hz, 1H), 3.15 (dd, *J*=11.7, 7.0 Hz, 1H), 1.45 (s, 3H), 1.43 (s, 9H), 1.41 (s, 3H) ppm.

B. *tert*-Butyl (1*R*)-1-{[(4*R*)-4-cyano-1,3-thiazolidin-3-yl]carbonyl}-2-[(4-methoxybenzyl)thio]-2-methylpropylcarbamate.

To a stirred solution of compound A (288 mg, 0.595 mmol) in CH₂Cl₂ (6 mL) was added trifluoroacetic anhydride (0.168 mL, 1.19 mmol). The reaction mixture was stirred at RT for 5 hours. The reaction mixture was concentrated *in vacuo* and purified via flash chromatography over silica gel (2:1 hexanes:EtOAc) to give 84 mg (30% yield) of compound B as a colorless oil.


¹H NMR (CDCl₃) 400 MHz δ 7.28 (d, *J*=8.6 Hz, 2H), 6.83 (d, *J*=8.6 Hz, 2H), 5.40 (d, *J*=8.9 Hz, 1H), 5.29 (dd, *J*=5.6, 3.7 Hz, 1H), 4.90-4.83 (m, 2H), 4.51 (d, *J*=9.2 Hz, 1H), 3.80 (s, 2H), 3.78 (s, 3H), 3.33-3.26 (m, 2H), 1.44 (s, 3H), 1.43 (s, 9H), 1.40 (s, 3H) ppm.

C. (4*R*)-3-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl}-1,3-thiazolidine-4-carbonitrile hydrochloride.

To a stirred solution of compound B (84 mg, 0.18 mmol) in 1,4-dioxane (1 mL) was added a solution of 4.0 M HCl in 1,4-dioxane (1.0 mL, 4.0 mmol). The reaction mixture was stirred at RT for 12 hours. The solvent was removed *in vacuo*, and the resulting oil was triturated with Et₂O to produce a light yellow solid. The solid was filtered *in vacuo*, washed with several portions of Et₂O, and dried *in vacuo* to produce 45 mg (62% yield) of the crude product. This material was purified via semi-preparative HPLC (10% acetonitrile in water ramped over 10 minutes to 90% acetonitrile in water) followed by re-salting with 2.0 M HCl in Et₂O to produce 5 mg (7% overall yield) of compound C as a light yellow solid.

¹H NMR (MeOH-d₄) 400 MHz δ 7.35 (d, *J*=8.6 Hz, 2H), 6.88 (d, *J*=8.6 Hz, 2H), 5.29 (t, *J*=5.5 Hz, 1H), 4.74 (d, *J*=9.2 Hz, 1H), 4.61 (d, *J*=9.2 Hz, 1H), 4.17 (s, 1H), 3.91 (dd, *J*=26.7, 12.7 Hz, 2H), 3.77 (s, 3H), 3.51-3.42 (m, 2H), 1.57 (s, 3H), 1.44 (s, 3H) ppm.

Example 2



(2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl}pyrrolidine-2-carbonitrile hydrochloride

A. *tert*-Butyl (1*R*)-1-{[(2*S*)-2-cyanopyrrolidin-1-yl]carbonyl}-2-[(4-methoxybenzyl)thio]-2-methylpropylcarbamate.

To a stirred solution of *N*-BOC-L-Pen(MOB)-OH (200 mg, 0.541 mmol) in DMF (6 mL) was added (2*S*)-pyrrolidine-2-carbonitrile 4-methylbenzenesulfonate (this compound was prepared as previously described in: *Bioorg. Med. Chem. Lett.* 1996, 6, 1163, Ashworth, D. M. et al., incorporated herein by reference in such regard (145 mg, 0.541 mmol), HATU (206 mg, 0.541 mmol), and *N,N*-diisopropylethylamine (0.380 mL, 2.164 mmol). The reaction mixture was stirred at RT for approximately 60 hours. Water (6 mL) was added and the reaction mixture was extracted with four portions of EtOAc. The combined extracts were washed with water, saturated CuSO₄, brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The crude material was flash chromatographed over silica gel (1:1 hexanes:EtOAc) to afford 213 mg (84% yield) of compound A as a colorless oil.

¹H NMR (CDCl₃) 400 MHz δ 7.27 (d, *J*=8.9 Hz, 2H), 6.82 (d, *J*=8.5 Hz, 2H), 5.44 (d, *J*=9.3 Hz, 1H), 4.82-4.79 (m, 1H), 4.52 (d, *J*=9.5 Hz, 1H), 3.91-3.83 (m, 2H), 3.80 (s, 2H), 3.77 (s, 3H), 2.29-2.10 (m, 4H), 1.43 (s, 12H), 1.40 (s, 3H) ppm.

B. (2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl}pyrrolidine-2-carbonitrile.

To a stirred solution of compound A (213 mg, 0.452 mmol) in CH₂Cl₂ (4 mL) was added TFA (1 mL). The reaction was stirred at RT for 4 hours. The solvent was removed *in vacuo* and the crude oil was re-dissolved in EtOAc and washed with sat. NaHCO₃. The aqueous layer was re-extracted with three portions of EtOAc. The combined extracts were dried over MgSO₄, decanted, and concentrated *in vacuo*. Purification via flash chromatography over silica gel (5% MeOH (with 2% NH₃) in CH₂Cl₂) afforded 75 mg (45% yield) of compound B as a white foam.


^1H NMR (CDCl_3) 400 MHz δ 7.27 (d, $J=8.4$ Hz, 2H), 6.83 (d, $J=8.6$ Hz, 2H), 4.77-4.75 (m, 1H), 3.77 (s, 3H), 3.74 (s, 2H), 3.63-3.46 (m, 3H), 2.26-2.07 (m, 4H), 1.91 (br s, 2H), 1.45 (s, 3H), 1.36 (s, 3H) ppm.

- 5 C. (2S)-1-[(2R)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl]pyrrolidine-2-carbonitrile hydrochloride.

Diethyl ether (4 mL) was added to a flask containing compound B (75 mg, 0.215 mmol). Several drops of acetone were added to allow the solution to become homogeneous. A solution of 2.0 M HCl in Et₂O (1.0 mL) was added and the reaction
10 mixture was stirred at RT for 5 minutes. A white solid precipitated during this time. The mixture was concentrated *in vacuo* to dryness and the solid was dried overnight under high vacuum to give 73 mg (88% yield) of compound C as a white solid.

^1H NMR (D_2O) 400 MHz δ 7.31 (d, $J=8.4$ Hz, 2H), 6.88 (d, $J=8.4$ Hz, 2H), 4.69-4.66 (m, 1H), 3.87-3.74 (m, 3H), 3.69 (s, 3H), 3.56-3.50 (m, 1H), 3.32-3.26 (m, 1H), 2.28-2.12
15 (m, 2H), 2.09-1.99 (m, 1H), 1.96-1.89 (m, 1H), 1.44 (s, 3H), 1.29 (s, 3H) ppm.

Example 3



- (2S)-1-[(2R)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl]pyrrolidine-
20 2-carbonitrile hydrochloride

A. *tert*-Butyl (1*R*)-1-{[(2*S*)-2-cyanopyrrolidine-1-yl]carbonyl}-2-[(4-methoxybenzyl)sulfonyl]-2-methylpropylcarbamate.

To a stirred solution of *tert*-butyl (1*R*)-1-{[(2*S*)-2-cyanopyrrolidin-1-yl]carbonyl}-2-[(4-methoxybenzyl)thio]-2-methylpropylcarbamate (427 mg, 0.905 mmol) in
5 chloroform (20 mL) at 0°C was added solid *m*-CPBA (1.56 g, 9.05 mmol) in one portion. The reaction mixture was stirred for 30 minutes at 0°C and then for 14 hours at RT. During this time the reaction mixture went from light purple to colorless to light yellow. The reaction mixture was washed with 1 M NaOH and separated. The aqueous layer was re-extracted with chloroform and the combined extracts were dried
10 over MgSO₄, filtered, and concentrated *in vacuo*. Purification via flash chromatography over silica gel (1:1 hexanes:EtOAc) afforded 355 mg (82% yield) of compound A as a white solid.

¹H NMR (CDCl₃) 400 MHz δ 7.32 (d, *J*=8.8 Hz, 2H), 6.88 (d, *J*=8.6 Hz, 2H), 5.49 (d, *J*=9.6 Hz, 1H), 5.15 (d, *J*=9.9 Hz, 1H), 4.76-4.73 (m, 1H), 4.22 (s, 2H), 3.93-3.88 (m, 1H),
15 3.82-3.78 (m, 1H), 3.78 (s, 3H), 2.29-2.12 (m, 4H), 1.55 (s, 3H), 1.47 (s, 3H), 1.42 (s, 9H) ppm.

B. (2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl}pyrrolidine-2-carbonitrile.

20 To a stirred solution of compound A (355 mg, 0.740 mmol) in CH₂Cl₂ (6.5 mL) was added TFA (1.5 mL). The reaction mixture was stirred at RT for 2 hours and then concentrated *in vacuo*. After re-dissolving in EtOAc, the reaction mixture was washed with saturated NaHCO₃. The aqueous layer was re-extracted with three portions of EtOAc, and the combined extracts were dried over MgSO₄, decanted, and concentrated


in vacuo. Purification via flash chromatography over silica gel (5% MeOH (with 2% NH₃) in CH₂Cl₂) afforded 74 mg (27% yield) of compound B as a light yellow oil.

¹H NMR (CDCl₃) 400 MHz δ 7.33 (d, *J*=8.8 Hz, 2H), 6.91 (d, *J*=8.8 Hz, 2H), 4.78-4.75 (m, 1H), 4.62 (d, *J*=13.4 Hz, 1H), 4.39 (s, 1H), 4.35 (d, *J*=13.3 Hz, 1H), 3.82-3.77 (m, 1H),
5 3.80 (s, 3H), 3.68-3.62 (m, 1H), 2.32-2.14 (m, 4H), 1.62 (s, 3H), 1.39 (s, 3H) ppm.

C. (2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl}pyrrolidine-2-carbonitrile hydrochloride.

Diethyl ether (4 mL) was added to a flask containing compound B (74 mg, 0.198
10 mmol). Several drops of acetone and CH₂Cl₂ were added to allow the solution to become homogeneous. A solution of 2.0 M HCl in Et₂O (2.0 mL) was added and the reaction mixture was stirred at RT for 5 minutes. A white solid precipitated during this time. The mixture was concentrated *in vacuo* to dryness and the solid was dried overnight under high vacuum to give 66 mg (80% yield) of compound C as a white
15 solid.

¹H NMR (D₂O) 400 MHz δ 7.28 (d, *J*=8.5 Hz, 2H), 6.93 (d, *J*=8.7 Hz, 2H), 4.75-4.71 (m, 2H), 4.53 (s, 2H), 3.70 (s, 3H), 3.66-3.59 (m, 2H), 2.27-2.17 (m, 2H), 2.09-1.94 (m, 2H), 1.63 (s, 3H), 1.47 (s, 3H) ppm.

Example 4

(4R)-3-[(2R)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl]-1,3-thiazolidine-4-carbonitrile 1,1-dioxide hydrochloride

- 5 A. *tert*-Butyl (1*R*)-1-{[(4*R*)-4-cyano-1,1-dioxido-1,3-thiazolidin-3-yl]carbonyl}-2-[(4-methoxybenzyl)sulfonyl]-2-methylpropylcarbamate.

To a stirred solution of *tert*-butyl (1*R*)-1-{[(4*R*)-4-cyano-1,3-thiazolidin-3-yl]carbonyl}-2-[(4-methoxybenzyl)thio]-2-methylpropylcarbamate (151 mg, 0.324 mmol) in chloroform (8 mL) was added *m*-CPBA (560 mg, 3.24 mmol). The reaction mixture was stirred at RT for 14 hours. The reaction mixture was then washed with 1 M NaOH and separated. The aqueous layer was re-extracted with chloroform and the combined extracts were dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification via flash chromatography over silica gel (1:1 hexanes:EtOAc) afforded 121 mg (70% yield) of compound A as a white solid.

- 15 ¹H NMR (CDCl₃) 400 MHz δ 7.33 (d, *J*=8.2 Hz, 2H), 6.92 (d, *J*=8.6 Hz, 2H), 5.50–5.42 (m, 2H), 5.21–5.12 (m, 1H), 4.88 (d, *J*=9.2 Hz, 1H), 4.71 (d, *J*=11.5 Hz, 1H), 4.29–4.20 m, 2H), 3.81 (s, 3H), 3.62–3.45 (m, 2H), 1.54 (s, 3H), 1.52 (s, 3H), 1.43 (s, 9H) ppm.

- 20 B. (4*R*)-3-[(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl]-1,3-thiazolidine-4-carbonitrile 1,1-dioxide.


To a stirred solution of compound A (121 mg, 0.228 mmol) in CH₂Cl₂ (2.0 mL) was added TFA (0.5 mL). The reaction mixture was stirred at RT for 2 hours and then concentrated *in vacuo*. After re-dissolving in EtOAc, the reaction mixture was washed with saturated NaHCO₃. The aqueous layer was re-extracted with three portions of EtOAc, and the combined extracts were dried over MgSO₄, decanted, and concentrated *in vacuo*. Purification via flash chromatography over silica gel (2% MeOH (with 2% NH₃) in CH₂Cl₂ to 5% MeOH (with 2% NH₃) in CH₂Cl₂) afforded 48 mg (49% yield) of compound B as a light yellow oil.

¹H NMR (CDCl₃) 400 MHz δ (major rotomer) 7.32 (d, *J*=8.4 Hz, 2H), 6.92 (d, *J*=8.6 Hz, 2H), 5.52 (dd, *J*=8.1, 5.2 Hz, 1H), 4.91 (d, *J*=11.6 Hz, 1H), 4.56 (d, *J*=9.3 Hz, 1H), 4.53 (d, *J*=11.1 Hz, 1H), 4.32-4.06 (m, 2H), 3.81 (s, 3H), 3.60-3.51 (m, 2H), 1.99 (br s, 2H), 1.57 (s, 3H), 1.40 (s, 3H) ppm.

C. (4*R*)-3-[(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl]-1,3-thiazolidine-4-carbonitrile 1,1-dioxide hydrochloride.

Diethyl ether (4 mL) was added to a flask containing compound B (48 mg, 0.112 mmol). Several drops of acetone were added to allow the solution to become homogeneous. A solution of 2.0 M HCl in Et₂O (1.0 mL) was added and the reaction mixture was stirred at RT for 5 minutes. A white solid precipitated during this time. The mixture was concentrated *in vacuo* to dryness and the solid was dried overnight under high vacuum to give 36 mg (86% yield) of compound C as a white solid.

¹H NMR (D₂O) 400 MHz δ 7.30 (d, *J*=8.0 Hz, 2H), 6.95 (d, *J*=8.0 Hz, 2H), 5.62-5.59 (m, 1H), 4.98 (d, *J*=11.5 Hz, 1H), 4.88 (d, *J*=11.3 Hz, 1H), 4.70 (s, 1H), 4.59-4.51 (m, 2H), 3.99-3.87 (m, 2H), 3.72 (s, 3H), 1.64 (s, 3H), 1.52 (s, 3H) ppm.

Example 5

5

(2S)-1-[(2S)-2-Amino-3,3-bis(4-fluorophenyl)propanoyl]pyrrolidine-2-carbonitrile hydrochloride.

A. tert-Butyl (1S)-1-[bis(4-fluorophenyl)methyl]-2-[(2S)-2-cyanopyrrolidin-1-yl]-2-oxoethylcarbamate.

10 To a stirred solution of (2S)-2-[(tert-butoxycarbonyl)amino]-3,3-bis(4-fluorophenyl)propanoic acid (475 mg, 1.26 mmol; for the preparation of this compound see below) in DMF (12 mL) was added (2S)-pyrrolidine-2-carbonitrile 4-methylbenzenesulfonate (338 mg, 1.26 mmol), HATU (479 mg, 1.26 mmol), and diisopropylethylamine (0.658 mL, 3.78 mmol). The reaction mixture was stirred at RT

15 for 16 hours then diluted with water (10 mL). The reaction mixture was extracted with 4 portions of EtOAc, and the combined extracts were washed with water, brine, dried over MgSO₄, filtered, and concentrated *in vacuo*. The crude product was flash chromatographed over silica gel (1:1 hexanes:EtOAc) to afford 335 mg (58% yield) of compound A as a white foam.

20 ¹H NMR (CDCl₃) 400 MHz δ 7.31-7.25 (m, 2H), 7.21-7.16 (m, 2H), 7.05-6.96 (m, 4H), 5.05-4.96 (m, 2H), 4.62 (t, J=5.5 Hz, 1H), 4.40 (d, J=11.0 Hz, 1H), 3.57 (q, J=9.0 Hz,

1H), 2.75-2.70 (m, 1H), 2.10-2.05 (m, 2H), 1.97-1.88 (m, 1H), 1.81-1.70 (m, 1H), 1.33 (s, 9H) ppm.

B. (2S)-1-[(2S)-2-Amino-3,3-bis(4-fluorophenyl)propanoyl]pyrrolidine-2-carbonitrile.

5 To a stirred solution of compound A (300 mg, 0.659 mmol) in CH₂Cl₂ (7 mL) was added TFA (0.507 mL, 6.59 mmol). The reaction mixture was stirred at RT for 12 hours followed by concentration *in vacuo*. The reaction mixture was re-dissolved in EtOAc and washed with saturated NaHCO₃, dried over MgSO₄, filtered, and concentrated *in vacuo*. Purification via flash chromatography over silica gel (3%
10 MeOH (with 2% NH₃) in CH₂Cl₂) afforded 129 mg (55% yield) of compound B as a white solid.

¹H NMR (CDCl₃) 400 MHz δ 7.33-7.30 (m, 2H), 7.22-7.19 (m, 2H), 7.09 (t, *J*=8.5 Hz, 2H), 6.98 (t, *J*=8.6 Hz, 2H), 4.65 (dd, *J*=7.7, 3.5 Hz, 1H), 4.30 (d, *J*=10.2 Hz, 1H), 4.09 (d, *J*=10.1 Hz, 1H), 3.37 (q, *J*=9.0 Hz, 1H), 2.67 (dt, *J*=8.6, 3.6 Hz, 1H), 2.22-2.00 (m, 4H),
15 1.96-1.87 (m, 1H), 1.81-1.70 (m, 1H) ppm.

C. (2S)-1-[(2S)-2-Amino-3,3-bis(4-fluorophenyl)propanoyl]pyrrolidine-2-carbonitrile hydrochloride.

Diethyl ether (6 mL) was added to a flask containing compound B (129 mg,
20 0.363 mmol). Several drops of acetone were added in order to allow the solution to become homogeneous. A solution of 2.0 M HCl in Et₂O (2.0 mL) was added and the reaction mixture was stirred at RT for 5 minutes. A white solid precipitated during this time. The solid was collected via vacuum filtration on a glass frit and dried

overnight under high vacuum to give 117 mg (82% yield) of compound C as a white solid.

^1H NMR (D_2O) 400 MHz δ 7.52-7.45 (m, 2H), 7.31-7.24 (m, 2H), 7.10 (t, $J=7.3$ Hz, 2H), 6.98 (t, $J=7.9$ Hz, 2H), 4.83 (d, $J=11.3$ Hz, 1H), 4.60-4.54 (m, 1H), 4.41 (d, $J=11.1$ Hz, 1H), 3.44-3.34 (m, 1H), 2.78-2.69 (m, 1H), 2.12-2.01 (m, 1H), 1.97-1.87 (m, 1H), 1.83-1.72 (m, 1H), 1.58-1.46 (m, 1H) ppm.

Preparation of (2S)-2-[(tert-butoxycarbonyl)amino]-3,3-bis(4-fluorophenyl)propanoic acid (Referenced above in Example 5)

10 A. 3,3-Bis(4-fluorophenyl)-3-hydroxypropanoic acid.

To an anhydrous THF (80 mL) solution of n-butyl lithium (46 mL of 2.5 M, 115 mmol) at 0 °C was added dropwise diisopropylamine (11.13 g, 115 mmol) and the solution stirred for 10 minutes. Keeping the solution at 0 °C, acetic acid (2.64 g, 44 mmol) was added dropwise and the mixture stirred for 10 min and it was then heated
15 50 °C. After 30 min a heavy precipitate had formed and the solution was allowed to cool. A solution of 4,4'-difluorobenzophenone (9.6 g, 0.044 mol) in THF (50 mL, anhydrous) was added at 0 °C, and the solution stirred at room temperature overnight. Water (100 mL) and diethyl ether (100 mL) were added and the aqueous layer was separated and acidified with 1M HCl to pH 3. The organics were extracted with ethyl
20 acetate (3 X 200 mL) followed by drying over MgSO_4 . Filtration and removal of the solvent *in vacuo* yielded a crude white solid that could be washed with cold CHCl_3 to remove trace amounts of the benzophenone. The solid was dried under high vacuum yielding 5.63 g (20.2 mmol, 46% yield) of compound A as a white solid.

^1H NMR (d_6 -DMSO) 400 MHz δ 12.4 (s(br), 1H), 7.48-7.39 (m, 4H), 7.19-7.02 (m, 4H), 5.91 (s(br), 1H), 3.25 (s, 2H) ppm.

B. 3,3-Bis(4-fluorophenyl)acrylic acid.

5 To a 20% solution of sulfuric acid in acetic acid (50 mL, V/V) was compound A (5.6 g, 20.2 mmol) and the mixture stirred for 30 minutes at RT. To this solution was added H_2O (500 mL) and the organics were extracted with ethyl acetate (3 X 150 mL) followed by drying over MgSO_4 . Filtration and removal of the solvent *in vacuo* yielded a white solid. The solid was dried under high vacuum yielding 4.97g (19.1mmol, 95%
10 yield) of compound B as a white solid.

^1H NMR (CDCl_3) 400 MHz δ 7.27-7.21 (m, 2H), 7.19-7.13 (m, 2H), 7.10-6.95 (m, 4H), 6.26 (s, 1H) ppm.

C. 3,3-Bis(4-fluorophenyl)propanoic acid.

15 To a solution of compound B (2.5 g, 9.61mmol) in ethyl acetate (250 mL) was added 10% palladium on carbon (50% w/w) and hydrogenated at 1 atmosphere of hydrogen for 12 hours. The heterogeneous solution was filtered through celite and concentrated *in vacuo* to provide a yellow oil. The oil was dried under high vacuum yielding 2.40 g (9.16 mmol, 95 % yield) of compound C as a yellow oil.

20 ^1H NMR (d_6 -DMSO) 400 MHz δ 12.08 (brs, 1H), 7.40-7.30 (m, 4H), 7.15-7.05 (m, 4H), 4.45 (t, 1H, $J = 8.1$ Hz), 3.05(d, 2H, $J = 8.1$ Hz) ppm.

D. (4*S*,5*R*)-3-[3,3-Bis(4-fluorophenyl)propanoyl]-4-methyl-5-phenyl-1,3-oxazolidin-2-one.

To a THF (50 mL, anhydrous) containing compound C (2.0 g, 7.63 mmol) was added N,N-diisopropylethylamine (1.18 g, 9.16 mmol) and then the solution cooled to -78 °C. To this solution was added trimethylacetyl chloride (0.97 g, 8.01 mmol) and the solution warmed to 0 °C over 1 hour. The cloudy mixture was filtered and the
5 filtrate added slowly over 10 min to a solution of the lithiated (4S,5R)-(-)-4-methyl-5-phenyl-2-oxazolidinone at -78 °C, which was prepared by the dropwise addition of n-butyl lithium (3.0 mL of 2.5 M, 7.63 mmol) to a THF (50 mL) solution of (4S,5R)-(-)-4-methyl-5-phenyl-2-oxazolidinone (1.35 g, 7.63 mmol) at -78 °C which had stirred for 10 min to provide the lithiated (4S,5R)-(-)-4-methyl-5-phenyl-2-oxazolidinone. The
10 yellow mixture was warmed to 0 °C and quenched with H₂O (50 mL) and extracted with diethyl ether (3 X 250 mL) followed by drying over MgSO₄. Filtration and removal of the solvent *in vacuo* yielded a solid. Flash chromatography (silica gel, 20 % ethyl acetate/ hexanes) provided compound D. The white solid was dried under high vacuum yielding 2.31g (5.49 mmol, 72% yield) as a white solid.

15 ¹H NMR (d₆-DMSO) 400 MHz δ 7.40-7.25 (m, 9H), 7.18-7.02 (m, 4H), 5.76 (d, 1H, *J* = 7.6 Hz), 4.65 (m, 1H), 4.58 (t, 1H, *J* = 7.6 Hz), 3.72 (dd, 1H, *J* = 16.8, 7.0 Hz) 3.57 (dd, 1H, *J* = 16.8, 7.0 Hz), 0.58 (d, 3H, *J* = 6.7 Hz) ppm.

E. (4S,5R)-3-[(2S)-2-Azido-3,3- bis(4-fluorophenyl) propanoyl]-4-methyl-5-[(1E,3Z)-
20 1-methylhexa-1,3,5-trienyl]-1,3-oxazolidin-2-one.

To a THF (50 mL, anhydrous) solution containing compound D (2.0 g, 4.75 mmol) at -78 °C was added dropwise potassium bis(trimethylsilyl)amide (10.0 mL of 0.5 M toluene solution, 4.98 mmol). After stirring for 10 min 2,4,6-triisopropylbenzenesulfonyl azide (trisyl azide) (1.84 g, 5.94 mmol) in THF (10 mL,

anhydrous) was added in one portion. After 3 minutes acetic acid was added (1.31 g, 21.8 mmol) at -78°C and then the reaction quickly warmed to 30°C and stirred for 1 hr at that temperature generating a light yellow solution. To this solution was added H_2O (100 mL) and the organics were extracted with ethyl acetate (500 mL). After
5 washing with sat. NaHCO_3 (100 mL) and drying over MgSO_4 the solvent was removed *in vacuo* yielding a yellow oil. Column chromatography (ethyl acetate/hexanes 1:9) provided compound E as a white solid. HPLC showed a single diastereoisomer. The white solid was dried under high vacuum yielding 1.71 g (3.70 mmol, 78% yield) as a white solid.

10 ^1H NMR (CDCl_3) 400 MHz δ 7.42-7.35 (m, H), 7.25-7.18 (m, H), 7.10-7.06 (m, 2H), 7.05-6.92 (m, 2H), 5.95 (d, 1H, $J = 10.8$ Hz), 5.05 (d, 1H, $J = 7.1$ Hz), 4.60 (d, 1H, $J = 10.8$ Hz), 4.38 (m, 1H), 0.95 (d, 3H, $J = 6.8$ Hz) ppm.

F. (2S)-2-Azido-3,3-bis(4-fluorophenyl)propanoic acid.

15 To a THF/ H_2O (4:1, 50 mL) solution of compound E (1.5 g, 3.25 mmol) at 0°C was added a solution of lithium hydroxide (0.272 g, 6.49 mmol) in hydrogen peroxide (1.50 mL of 30% soln in H_2O , 48.75 mmol). The mixture was stirred at 0°C for 1 hr and then quenched with Na_2SO_4 (6.3 g, 50 mL of 1.0 M solution in H_2O). The THF was removed *in vacuo* and the solution acidified to pH 1 with 6.0 M HCl at 0°C . The
20 organics were extracted with ethyl acetate (2 X 200 mL) followed by drying over MgSO_4 . Filtration and removal of the solvent *in vacuo* yielded a clear oil. Column chromatography (EtOAc/hexanes/acetic acid 50:50:1) provided compound F as a white solid. The solid was dried under high vacuum yielding 0.78 g (2.60 mmol, 80% yield) as a white solid.

¹H NMR (CDCl₃) 400 MHz δ 9.60(s(br), 1H), 7.25-7.10 (m, 4H), 7.10-6.95 (m, 4H), 4.50 (d, 2H, *J* = 8.6 Hz) ppm.

G. (2S)-2-Amino-3,3-bis(4-fluorophenyl)propanoic acid.

5 To an ethyl acetate (250 mL) solution of compound F (1.5 g, 4.95 mmol) was added 10% palladium on carbon (10% w/w) and hydrogenated at 1 atmosphere of hydrogen for 12 hr. The heterogeneous solution was filtered through celite (1 g) and the filtrate concentrated *in vacuo* to provide a clear oil. The oil was dried under high vacuum yielding 1.30 g (4.70 mmol, 95% yield) of compound G as a white solid.

10 ¹H NMR (d₆-DMSO) 400 MHz δ 10.2(s(br), 1H), 7.38-7.27(m, 4H), 7.08-6.98 (m, 4H), 4.25 (d, 1H, *J* = 8.3 Hz), 3.95 (d, 1H, *J* = 8.3 Hz) ppm.

H. (2S)-2-[(*tert*-Butoxycarbonyl)amino]-3,3-bis(4-fluorophenyl)propanoic acid.

To a CH₂Cl₂ (150 mL) solution containing compound G (1.30 g, 4.69 mmol) was
15 added triethylamine (2.37 g, 23.4 mmol) and di-*tert*-butyl dicarbonate (1.23 g, 5.63 mmol). After stirring for 12 hr H₂O (50 mL) and CH₂Cl₂ (300 mL) were added and the solution acidified to pH 3 with 1.0 M HCl. Separation of the ethyl acetate layer followed by drying over MgSO₄ and removal of the solvent *in vacuo* yielded a clear oil. The oil was dried under high vacuum yielding 1.68 g (4.4 mmol, 95% yield) of
20 compound H as a white solid.

¹H NMR (d₆-DMSO) 400 MHz δ 12.4 (s(br), 1H), 7.35-7.22 (m, 4H), 7.15-6.95 (m, 4H), 4.78 (t, 1H, *J* = 8.9 Hz), 4.25 (d, 1H, *J* = 8.9 Hz), 3.05 (m, 1H), 1.20 (s, 3H), 1.15 (s, 6H) ppm.

BIOLOGICAL DATAMaterials:

H-Ala-Pro-pNA•HCl was purchased from BACHEM Bioscience Inc. (product no. L-1115). A 500 mM stock solution was prepared with dimethylsulfoxide and stored at
5 -20 °C. Gly-Pro-AMC was purchased from Enzyme System Products (product no. AMC-39) and stored at -20 °C as a 10 mM stock solution in dimethylsulfoxide. Test compounds were dissolved to 10 mM in dimethylsulfoxide and this was used as a stock solution for DPP-IV titration assays. Athens Research and Technology, Inc prepared the purified human DPP-IV. The material was isolated from human prostasomes using
10 the method of DeMeester et al., *J. Immunol. Methods* 189, 99-105. (1996), incorporated herein by reference in such regard.

DPP-IV Assay:

Two-fold serial dilutions of test compounds in 100 % dimethylsulfoxide were performed in 96-well polystyrene flat bottom plates (Costar, #9017). The average
15 enzymatic activity from wells containing dimethylsulfoxide but lacking test compound was used as a control value for calculating percent inhibition. DPP-IV (20 ng/mL) was mixed in microtiter plates with test compounds, substrate and assay buffer to yield 100 µM H-Ala-Pro-pNA•HCl in 25 mM Tris, pH 7.5, 10 mM KCl, 140 mM NaCl. The intact peptide contains a p-nitrophenylanilide which, when hydrolyzed by DPP-IV,
20 releases the absorbant p-nitrophenylaniline. The absorbency was monitored in 20 minutes intervals at a wavelength of 387 nm using a Molecular Devices SpectraMax 250 absorbency plate reader. The enzymatic activity was determined by estimating the best linear fit to the data. Values for enzymatic activity were taken directly from the linear fit determined by the software on the plate reader.

Data Analysis: The enzymatic activity was determined by estimating the best linear fit to the data. Data reduction was performed using the Microsoft Excel RoboSage.

Determination of IC₅₀ values: The enzymatic activity was plotted against the concentration of test compound, including [I] = 0, and the IC₅₀ determined from a fit of equation 2 to the data.

$$\text{RATE} = V_{\max} / (1 + ([I] / IC_{50})) \quad (2)$$

V_{max} was the best fit estimate of the maximal enzymatic activity.

Determination of K_i values: K_i values were calculated from IC₅₀ values using equation 3 assuming a competitive model.

$$K_i = IC_{50} * [1 - \frac{S}{(S + K_m)}] \quad (3)$$

The apparent pK_i values were > 5.0 for each of the examples.

DPP-II Assay:

The intermediate plate contained 5.3 µL of test compound in 2-fold serial dilutions across the plate. A volume of 209 µL of buffer (100 mM sodium acetate pH 5.5) containing substrate (H-Lys-Ala-pNA-2HCl; product no. L-2085; BACHEM Bioscience Inc.) was added to each well of the intermediate plate, then mixed. The reaction was initiated with the transfer of 180 µL of the substrate/test compound solution to the assay plate containing 20 µL of enzyme. Final concentrations in the assay were 100 nM enzyme and 1000 µM substrate in 100 mM NaOAc, pH 5.5, 2.5% DMSO in a final volume of 200 µL. The absorbance was monitored every 20 minutes

for 5 hours at 387 nm using a Molecular Devices SpectraMax 250 absorbance plate reader.

Data Analysis: The enzymatic activity was determined by estimating the best linear fit to the data. Data reduction was performed using the Microsoft Excel

5 RoboSage.

Determination of IC₅₀ values: The enzymatic activity was plotted against the concentration of test compound, including [I] = 0, and the IC₅₀ determined from a fit of equation 2 to the data.

$$\text{RATE} = V_{\max} / (1 + ([I] / IC_{50})) \quad (2)$$

10 V_{\max} was the best fit estimate of the maximal enzymatic activity.

Determination of K_i values: K_i values were calculated from IC₅₀ values using equation 3 assuming a competitive model.

$$K_i = IC_{50} * [1 - \frac{S}{(S + K_m)}] \quad (3)$$

Certain compounds of the present invention showed activity for DPP-II, for
15 example pK_i values of >7.0 were observed, while others demonstrated selectivity for DPP-IV, discussed hereinabove.

In vivo studies:

Age and weight matched male CD1 mice were housed individually at 72° F and
50% relative humidity with a 12 h light/dark cycle. Animals were dosed by oral
20 gavage with 10 ml/kg vehicle (0.5% methylcellulose (HPMC) with 0.1% Tween 80) or 1
mg/kg test compound in vehicle. The animals were anesthetized with isoflurane for


blood collection at the specified times (0-6 hours). Plasma DPP-IV activity was measured using the fluorogenic substrate Gly-Pro-AMC (50 μ M) according to the manufacturers specification (Enzyme System Products, Livermore CA). The substrate was mixed with 50 mM Tris, pH 7.8 and 20% plasma. The samples were incubated for
5 20 min at 30°C and fluorescence measured using a cytofluor spectrofluoremeter with the filters set at 360 nm excitation and 460 nm emission.

All research complied with the principles of laboratory animal care (NIH publication No. 85-23, revised 1985) and GlaxoSmithKline policy on animal use.

10 Although specific embodiments of the present invention have been illustrated and described in detail, the invention is not limited thereto. The above detailed description of preferred embodiments is provided for example only and should not be construed as constituting any limitation of the invention. Modifications will be obvious to those skilled in the art, and all modifications that do not depart from the
15 spirit of the invention are intended to be included within the scope of the appended claims.

What is claimed is:

1. A compound of formula (I):



including salts, solvates, and pharmaceutically functional derivatives thereof, wherein

X is $-S(O)_b-$ or $-CH_2-$;


b is 0-2;

R^1 and R^2 each are either

- (i) H;
- (ii) alkyl;
- (iii) optionally substituted aryl or heteroaryl; or
- (iv) combine to form a 3 to 14 membered ring system, optionally containing one or more heteroatoms, and optionally containing one or more degrees of unsaturation;

when R^1 and R^2 are (iii), then R^3 is H or alkyl;

when R^1 and R^2 are (i), (ii), or (iv), then R^3 is



where n is 0-5;

m is 0 to 12;


Y is $S(O)_p$, O, alkylene, alkenylene, alkynylene, or a bond;

p is 0-2;

R^4 is R^5 when Y is S, O, alkylene, alkenylene, alkynylene, or a bond, where R^5 is alkyl, alkenyl, alkynyl, aryl, cycloalkyl, heteroaryl, or heterocyclyl;

R^4 is R^6 when Y is $S(O)$ or $S(O)_2$, where R^6 is alkyl, aryl, cycloalkyl, heteroaryl, amino, alkylamino, arylamino, heteroaryl amino, or cycloalkyl amino.

2. The compound of claim 1 wherein R^1 and R^2 are each alkyl and R^3 is



3. The compound of claim 2 wherein each of said alkyl is C_1 - C_6 alkyl.
4. The compound of claim 3 wherein each of said alkyl is methyl.
5. The compound of claim 4 wherein X is $-S(O)_b$ -, b is 0, n is 0, m is 1, Y is $-S(O)_p$ -, p is 0, R^4 is R^5 , and R^5 is optionally substituted aryl.
6. The compound of claim 5 wherein R^5 is phenyl substituted with alkoxy.
7. The compound of claim 4 wherein X is $-CH_2$ -, n is 0, m is 1, Y is $-S(O)_p$ -, p is 0, R^4 is R^5 , and R^5 is optionally substituted aryl.
8. The compound of claim 7 wherein R^5 is phenyl substituted with alkoxy.
9. The compound of claim 4 wherein X is $-CH_2$ -, n is 0, m is 1, Y is $S(O)_p$, p is 2, R^4 is R^6 , and R^6 is optionally substituted aryl.
10. The compound of claim 9 wherein R^6 is phenyl substituted with alkoxy.
11. The compound of claim 4 wherein X is $S(O)_b$, b is 2, n is 0, m is 1, Y is $S(O)_p$, p is 2, R^4 is R^6 , and R^6 is optionally substituted aryl.
12. The compound of claim 11 wherein R^6 is phenyl substituted with alkoxy.

13. The compound of claim 1 wherein the depicted NH₂ group is *cis* to the depicted nitrile warhead.
14. The compound of claim 1 wherein the depicted NH₂ group is *trans* to the depicted nitrile warhead.
15. A compound selected from:
 - (4*R*)-3-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl}-1,3-thiazolidine-4-carbonitrile hydrochloride;
 - (2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)thio]-3-methylbutanoyl}pyrrolidine-2-carbonitrile hydrochloride;
 - (2*S*)-1-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl}pyrrolidine-2-carbonitrile hydrochloride;
 - (4*R*)-3-{(2*R*)-2-Amino-3-[(4-methoxybenzyl)sulfonyl]-3-methylbutanoyl}-1,3-thiazolidine-4-carbonitrile 1,1-dioxide hydrochloride; and
 - (2*S*)-1-[(2*S*)-2-Amino-3,3-bis(4-fluorophenyl)propanoyl]pyrrolidine-2-carbonitrile hydrochloride.
16. A pharmaceutical formulation comprising a compound of claims 1 to 15.
17. The pharmaceutical formulation of claim 16 further comprising a pharmaceutically acceptable carrier.
18. A method of inhibiting a post proline/aniline cleaving protease comprising administering a compound of claims 1 to 15.
19. The method of claim 18 wherein the post proline/aniline cleaving protease is a serine protease.
20. The method of claim 19 wherein the serine protease is a dipeptidyl peptidase.
21. The method of claim 20 wherein the dipeptidyl peptidase is DPP-II.
22. The method of claim 20 wherein the dipeptidyl peptidase is DPP-IV.

23. A method for the treatment or prophylaxis of metabolic disorders, gastrointestinal disorders, viral disorders, inflammatory disorders, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders, encephalomyelitis, complement mediated disorders, glomerulonephritis, lipodystrophy, tissue damage, psychosomatic, depressive, and neuropsychiatric disorders, HIV infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions comprising administering a compound of claims 1 to 15.
24. The method of claim 23 wherein the compound of claims 1 to 15 is administered for the treatment or prophylaxis of diabetes.
25. Use of a compound of claims 1 to 15 in the manufacture of a medicament for the inhibition of a post proline/aniline cleaving protease.
26. The use of claim 25 wherein the post proline/aniline cleaving protease is a serine protease.
27. The use of claim 26 wherein the serine protease is a dipeptidyl peptidase.
28. The use of claim 27 wherein the dipeptidyl peptidase is DPP-II.
29. The use of claim 27 wherein the dipeptidyl peptidase is DPP-IV.
30. Use of a compound of claims 1 to 15 in the manufacture of a medicament for the treatment or prophylaxis of metabolic disorders, gastrointestinal disorders, viral disorders, inflammatory disorders, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders, encephalomyelitis, complement mediated disorders, glomerulonephritis, lipodystrophy, tissue damage, psychosomatic, depressive, and

neuropsychiatric disorders, HIV infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions.

31. A compound according to claims 1 to 15 for use as an active therapeutic substance.
32. A compound according to claims 1 to 15 for use in the manufacture of a medicament for the inhibition of serine protease.
33. A compound according to claims 1 to 15 for use in the manufacture of a medicament for the treatment or prophylaxis of metabolic disorders, gastrointestinal disorders, viral disorders, inflammatory disorders, diabetes, obesity, hyperlipidemia, dermatological or mucous membrane disorders, psoriasis, intestinal distress, constipation, autoimmune disorders, encephalomyelitis, complement mediated disorders, glomerulonephritis, lipodystrophy, tissue damage, psychosomatic, depressive, and neuropsychiatric disorders, HIV infection, allergies, inflammation, arthritis, transplant rejection, high blood pressure, congestive heart failure, tumors, and stress-induced abortions.